

Chemistry 12

Equilibrium Review Package



Name:

Date:

Block:

I. Multiple Choice:

1. Consider the rate diagram below for the following reaction:



Which of the following occurs at time t_1 ?

- A. addition of H_2
- B. addition of HI

- C. addition of a catalyst
- D. a decrease in volume

2. Chemical equilibrium is said to be dynamic because:


- A. the reaction proceeds quickly.
- B. the mass of the reactants is decreasing.
- C. the macroscopic properties are constant.
- D. both forward and reverse reactions are occurring.

3. Which reaction characteristics are changed by the addition of a catalyst to a reaction at constant temperature?

- I. Activation energy
- II. Equilibrium concentrations
- III. Reaction enthalpy

- A. I only
- B. III only
- C. I and II only
- D. I, II and III

4. Given the following system:



Which of the following chemicals, when added to the above system at equilibrium, would result in a decrease in $[\text{CrO}_4^{2-}]$?


- A. NaOH
- B. HNO_3

- C. Na_2CrO_4
- D. $\text{Na}_2\text{Cr}_2\text{O}_7$

5. Addition of a catalyst to an equilibrium system:

- A. increases the value of K_{eq}
- B. increases the yield of products.
- C. has no effect on the rates of reaction.
- D. increases the rate of formation of both reactants and product

6. Consider the following reaction:



The equilibrium expression is:

A. $K_{eq} = \frac{[2\text{BF}_3]}{[3\text{F}_2]}$

B. $K_{eq} = \frac{[\text{F}_2]^3}{[\text{BF}_3]^2}$


C. $K_{eq} = \frac{[\text{BF}_3]^2}{[\text{F}_2]^3}$

D. $K_{eq} = \frac{[\text{BF}_3]^2}{[\text{B}]^2[\text{F}_2]^3}$

7. The value of K_{eq} can be changed only by:

- A. adding a catalyst.
- B. changing the temperature.
- C. changing the reactant concentration.
- D. changing the volume of the container.


8. Consider the following equilibrium:



When 0.40 mol of PCl_3 and 0.40 mol of Cl_2 are placed in a 1.00 L container and allowed to reach equilibrium, 0.244 mol of PCl_5 are present. From this information, the value of K_{eq} is

- A. 0.10 B. 0.30 C. 3.3 D. 10


9. Consider the following equilibrium:



A 1.0 L container is filled with 0.05 mol PCl_5 , 1.0 mol PCl_3 , and 1.0 mol Cl_2 . The system proceeds to the:

- A. left because $\text{Trial } K_{eq} > K_{eq}$
- B. left because $\text{Trial } K_{eq} < K_{eq}$
- C. right because $\text{Trial } K_{eq} > K_{eq}$
- D. right because $\text{Trial } K_{eq} < K_{eq}$

10. A sample of $\text{X}_{(g)}$ is placed in a vessel and brought to equilibrium according to the reaction:




When the temperature is decreased, the concentration of Y in the reaction vessel increases.

Which of the following could explain this observation?

- A. The molecules are colliding with less energy, so the reaction shifts to the right.
- B. The reaction is endothermic, so when the temperature is decreased, the reaction shifts to the right.
- C. When the temperature is decreased, the value of K_{eq} increases, so the reaction shifts to the right.
- D. When the temperature is decreased, the value of $\text{Trial } K_{eq}$ also decreases, so the reaction shifts to the right.


11. Consider the following equilibrium:



Nitrogen gas and oxygen gas react when placed in a closed container. As the reaction proceeds towards equilibrium, the rate of the reverse reaction:

- A. increases as the [] of products decreases.
- B. decreases as the [] of products decreases.
- C. increases as the [] of products increases.
- D. decreases as the [] of products increases.


12. Consider the following equilibrium:



For the above reaction:

- A. both min. enthalpy and max. entropy favour products.
- B. both min. enthalpy and max. entropy favour reactants.
- C. min. enthalpy favours reactants and max. entropy favours products.
- D. min. enthalpy favours products and max. entropy favours reactants.


13. Consider the following equilibrium:



At constant temperature and volume, more I_2 is added to the above equilibrium. A new state of equilibrium results from a shift to the:

- A. left with a net decrease in $[\text{H}_2]$.
- B. left with a net increase in $[\text{H}_2]$.
- C. right with a net increase in $[\text{H}_2]$.
- D. right with a net decrease in $[\text{H}_2]$.


14. Consider the following equilibrium:



In a 1.0 L container at equilibrium there are 1.0 mol NOCl, 0.70 mol NO and 0.40 mol Cl_2 . At constant temperature and volume, 0.10 mol NOCl is added. The concentrations in the "new" equilibrium in comparison to the concentrations in the "old" equilibrium are:

	$[\text{NOCl}]$	$[\text{NO}]$	$[\text{Cl}_2]$
A.	new = old	new = old	new = old
B.	new > old	new > old	new > old
C.	new < old	new < old	new > old
D.	new < old	new > old	new > old

15. Consider the following equilibrium:



When 0.1010 mol H_2O is placed in a 1.000 L container, equilibrium is established. The equilibrium concentration of O_2 is 0.0010M. The equilibrium concentrations of H_2O and H_2 are:

	$[\text{H}_2\text{O}]$	$[\text{H}_2]$
A.	0.0990	0.0020
B.	0.1000	0.0010
C.	0.1005	0.0005
D.	0.1010	0.0020

16. Consider the following equilibrium: $2\text{CO}_{(g)} + \text{O}_{2(g)} \rightleftharpoons 2\text{CO}_{2(g)}$

The ratio used to calculate the equilibrium constant is:


A. $\frac{[\text{2CO}]^2[\text{O}_2]}{[\text{2CO}_2]^2}$

B. $\frac{[\text{2CO}_2]^2}{[\text{2CO}]^2[\text{O}_2]}$

C. $\frac{[\text{CO}]^2[\text{O}_2]}{[\text{CO}_2]^2}$

D. $\frac{[\text{CO}_2]^2}{[\text{CO}]^2[\text{O}_2]}$


17. Consider the following equilibrium:



A change in temperature of the above system increases the value of the equilibrium constant. The new state of equilibrium was established by a shift:

- A. left as a result of a decrease in temperature.
- B. right as a result of a decrease in temperature.
- C. left as a result of an increase in temperature.
- D. right as a result of an increase in temperature.


18. Consider the following equilibrium:



In a 1.0 L container at equilibrium there are 0.050 mol H_2 , 0.050 mol S and 1.0 mol H_2S . The value of K_{eq} is:

- A. 2.5×10^{-3} B. 5.0×10^{-2} C. 2.0×10^1 D. 4.0×10^2


19. Consider the following equilibrium:



At high temperature, H_2O and CO are placed in a closed container. As the system approaches equilibrium, the:

- A. rate of the forward and reverse reactions both increase.
- B. rate of the forward and reverse reactions both decrease.
- C. rate of the forward reaction decreases and the rate of the reverse reaction increases.
- D. rate of the forward reaction increases and the rate of the reverse reaction decreases.

20. Consider the following equilibrium:



The equilibrium shifts right when:

- A. NO_2 is added.
- B. N_2O_4 is removed.
- C. the temperature is decreased.
- D. the volume of the system is increased.


21. In an endothermic equilibrium system, the:

- A. minimum enthalpy and the maximum entropy both favour products.
- B. minimum enthalpy and the maximum entropy both favour reactants.
- C. minimum enthalpy favours products and the maximum entropy favours reactants.
- D. minimum enthalpy favours reactants and the maximum entropy favours products.

22. An equilibrium system shifts left when the temperature is increased. The forward reaction is

- A. exothermic and ΔH is positive.
- B. exothermic and ΔH is negative.
- C. endothermic and ΔH is positive.
- D. endothermic and ΔH is negative.

23. Given the following equilibrium system:



The equilibrium constant expression for the above system is:

- A. $K_{eq} = \frac{[\text{Br}_2(\text{l})]}{[\text{Br}_2(\text{g})]}$
- B. $K_{eq} = [\text{Br}_2(\text{g})]$
- C. $K_{eq} = \frac{1}{[\text{Br}_2(\text{g})]}$
- D. $K_{eq} = [\text{Br}_2(\text{g})][\text{Br}_2(\text{g})]$

24. Consider the following equilibrium:



When the temperature decreases, the equilibrium:

- A. shifts left and K_{eq} value increases.
- B. shifts left and K_{eq} value decreases.
- C. shifts right and K_{eq} value increases.
- D. shifts right and K_{eq} value decreases.

25. Consider the following equilibrium: $\text{CH}_4(\text{g}) + \text{H}_2\text{O}(\text{g}) \rightleftharpoons \text{CO}(\text{g}) + 3\text{H}_2(\text{g})$ $K_{eq} = 5.7$

At equilibrium, the $[\text{CH}_4] = 0.40\text{M}$, $[\text{CO}] = 0.30\text{M}$ and $[\text{H}_2] = 0.80\text{M}$. The $[\text{H}_2\text{O}] =$ is:

- A. 0.067M
- B. 0.11M
- C. 2.2M
- D. 5.3M

26. Consider the following equilibrium:



If 0.060 mol of O_3 and 0.70 mol of O_2 are introduced into a 1.0 L vessel, the

- A. Trial $K_{eq} > K_{eq}$ and the $[\text{O}_2]$ increases.
- B. Trial $K_{eq} < K_{eq}$ and the $[\text{O}_2]$ increases.
- C. Trial $K_{eq} > K_{eq}$ and the $[\text{O}_2]$ decreases.
- D. Trial $K_{eq} < K_{eq}$ and the $[\text{O}_2]$ decreases.


27. Macroscopic properties become constant in an equilibrium system when:

- A. all reactions have stopped.
- B. the reactants are completely used up.
- C. maximum enthalpy has been reached.
- D. forward and reverse reaction rates are equal.


28. In which of the following systems would the tendencies toward minimum enthalpy and maximum entropy be in opposition to each other?

- A. $\text{Br}_2(l) + \text{heat} \rightleftharpoons \text{Br}_2(g)$
- B. $\text{NaOH}(s) \rightleftharpoons \text{Na}^+(aq) + \text{OH}^-(aq) + \text{heat}$
- C. $2\text{C}(g) + 2\text{H}_2(g) \rightleftharpoons \text{C}_2\text{H}_4(g)$ ΔH is positive
- D. $2\text{K}(s) + 2\text{H}_2\text{O}(l) \rightleftharpoons 2\text{K}^+(aq) + 2\text{OH}^-(aq) + \text{H}_2(g)$ ΔH is negative

29. An uncatalyzed reaction was found to produce 40 kJ of energy in 10 minutes. When catalyzed, the same reaction produced 40 kJ of energy in 2 minutes. Which one of the following potential energy diagrams is consistent with the above data?




30. Consider the following equilibrium system:



Which one of the following statements describes the effect that a decrease in volume would have on the position of equilibrium and the $[\text{H}_2]$ in the above system?

- A. No shift, $[\text{H}_2]$ increases.
- B. Shift right, $[\text{H}_2]$ increases.
- C. Shift right, $[\text{H}_2]$ decreases.
- D. No shift, $[\text{H}_2]$ remains constant.


31. Tooth enamel, $\text{Ca}_5(\text{PO}_4)_3\text{OH}$ establishes the following equilibrium:



Which one of the following, when added to the above equilibrium system, would result in a shift to the right?

- A. $\text{H}^+(aq)$
- B. $\text{OH}^-(aq)$
- C. $\text{Ca}^{2+}(aq)$
- D. $\text{Ca}_5(\text{PO}_4)_3\text{OH}(s)$

37. Consider the following equilibrium:



Which of the following factors will not alter the position of equilibrium?

- A. a pressure decrease
 B. a temperature increase
 C. the presence of a catalyst
 D. the addition of more $\text{N}_2(\text{g})$


38. Which of the following is **least** likely to favour the formation of products?

- A. $2\text{H}_2\text{O}(\text{g}) \rightleftharpoons 2\text{H}_2(\text{g}) + \text{O}_2(\text{g})$ $K_{\text{eq}} = 7.3 \times 10^{-18}$
 B. $\text{N}_2\text{O}(\text{g}) + \text{NO}_2(\text{g}) \rightleftharpoons 3\text{NO}(\text{g})$ $K_{\text{eq}} = 4.2 \times 10^{-4}$
 C. $\text{N}_2\text{O}_4(\text{g}) \rightleftharpoons 2\text{NO}_2(\text{g})$ $K_{\text{eq}} = 4.5$
 D. $\text{SO}_2(\text{g}) + \text{NO}_2(\text{g}) \rightleftharpoons \text{NO}(\text{g}) + \text{SO}_3(\text{g})$ $K_{\text{eq}} = 85$

39. Consider the following equilibrium:



Which of the following graphs shows the result of increasing the temperature at time t_1 ?



40. Consider the following equilibrium and the table of experimental data:



	Initial		Equilibrium	
	$[\text{N}_2\text{O}_4]$	$[\text{NO}_2]$	$[\text{N}_2\text{O}_4]$	$[\text{NO}_2]$
Trial 1	0.0400	0.0000	0.0337	0.0125
Trial 2	0.0200	0.0600	0.0429	0.0141

Which of the following represents the K_{eq} value?

- A. 4.64×10^{-3}
 B. 3.71×10^{-1}
 C. 7.42×10^{-1}
 D. 2.16×10^2


II. Short Answers:

1. Consider the following equilibrium system:



A closed flask is found to contain 0.40M $\text{NO}_{(g)}$, 0.32M $\text{Cl}_{2(g)}$ and 5.6M $\text{NOCl}_{(g)}$. Determine the direction the reaction proceeds to reach equilibrium.

2. Consider the following equilibrium system:




The system is said to "shift right" as the result of the addition of **extra** $\text{H}_{2(g)}$. Describe the sequence of changes in both forward and reverse reaction rates as the system goes from the original equilibrium to the new equilibrium.

3. Consider the following equilibrium system:




A 2.00L container is filled with 0.500 mol of COF_2 . Calculate the $[\text{COF}_2]$ at equilibrium.

4. Consider the following equilibrium system:




Cooling the equilibrium changes the colour from green to blue. What effect will the decrease in temperature have on K_{eq} ? Explain, using Le Chatelier's Principle.

5. Consider the following:




Initially, 0.0600 mol CH₄, 0.0800 mol H₂O, 0.280 mol CO and 0.740 mol H₂ are placed into a 4.00 L container. At equilibrium, the [H₂] = 0.200M. What is the value of K_{eq}?

6. Consider the following equilibrium:



A 2.0 L flask is filled with 0.10 mol HI . Calculate the concentration of H₂ at equilibrium.

7. A flask is initially filled with some HF. At equilibrium, the [HF]=0.80M. What is the [H₂] at equilibrium?



Answers:

I. Multiple Choice:

- | | | | |
|-------|-------|-------|-------|
| 1) B | 11) C | 21) D | 31) A |
| 2) D | 12) D | 22) B | 32) B |
| 3) A | 13) A | 23) C | 33) C |
| 4) B | 14) B | 24) B | 34) A |
| 5) D | 15) A | 25) A | 35) A |
| 6) C | 16) D | 26) C | 36) C |
| 7) B | 17) B | 27) D | 37) C |
| 8) D | 18) D | 28) A | 38) A |
| 9) A | 19) C | 29) A | 39) A |
| 10) C | 20) D | 30) A | 40) A |

II. Short Answers:

- 1) Equilibrium shifts left. Trial $K_{eq} = 6.1 \times 10^2$
- 2) When conc. of H_2 increases, rate forward increases. As time proceeds, more HI is produced, therefore, rate reverse will increase. At equilibrium, both rate forward and reverse are constant.
- 3) $[COF_2] = 0.065M$
- 4) K_{eq} decreases
- 5) $K_{eq} = 4.00$
- 6) $[H_2] = 0.024M$
- 7) $[H_2] = 0.40M$